Contents Volume 63, 1988

Overview: ecological and evolutionary implications of taphonomic processes
S. M. Kidwell (Chicago, IL, U.S.A.) and A. K. Behrensmeyer (Washington, DC, U.S.A.) 1
Nautiloid shell taphonomy: interpretations based on water pressure
R. A. Hewitt (Hamilton, Ont., Canada) 15
Fossilization potential of the mud crab, *Panopeus* (Brachyura: Xanthidae) and temporal variability in crustacean taphonomy
R. E. Plotnick (Chicago, IL, U.S.A.), T. Baumiller (Chicago, IL, U.S.A.) and K. L. Wetmore (Baltimore, MD, U.S.A.) .. 27
Taphonomic significance of hermit crabs (Anomura: Paguridea): Epifaunal hermit crab—infaunal gastropod example
S. E. Walker (Berkeley, CA, U.S.A.) 45
The paleoecological significance of diversity: the effect of time averaging and differential preservation on macroinvertebrate species richness in death assemblages
G. M. Staff and E. N. Powell (College Station, TX, U.S.A.) 73
Taphonomy and preservation of a monospecific titanothere assemblage from the Washakie Formation (Late Eocene), southern Wyoming. An ecological accident in the fossil record
W. D. Turnbull (Chicago IL, U.S.A.) and D. M. Martill (Leicester, U.K.) 91
Taphonomy and depositional dynamics of Devonian shell-rich mudstones
K. M. Parsons, C. E. Brett and K. B. Miller (Rochester, NY, U.S.A.) 109
Sampling and faunal turnover in Early Eocene mammals
C. Badgley and P. D. Gingerich (Ann Arbor, MI, U.S.A.) 141
Paleoenvironmental controls on vertebrate-bearing abandoned channels in the Upper Carboniferous
R. W. Hook (Washington, DC, U.S.A.) and J. C. Ferm (Lexington, KY, U.S.A.) 159
Vertebrate preservation in fluvial channels
A. K. Behrensmeyer (Washington, DC, U.S.A.) 183
Taphonomic comparison of passive and active continental margins: Neogene shellbeds of the Atlantic Coastal Plain and Northern Gulf of California
S. M. Kidwell (Chicago, IL, U.S.A.) 201
Taphofacies models for epeiric sea environments: Middle Paleozoic examples
S. E. Speyer (Edmonton, Alta., Canada) and C. E. Brett (Rochester, NY, U.S.A.) 225
Taphonomic bias in fish diversity from Cenozoic floodplain environments
G. R. Smith, R. F. Stearly and C. E. Badgley (Ann Arbor, MI, U.S.A.) 263
The role of nutrient availability in bioerosion: consequences to carbonate buildups
P. Hallock (St. Petersburg, FL, U.S.A.) 275
Late Quaternary land snails from the north coast of Jamaica: local extinctions and climatic change
G. A. Goodfriend (Rehovot, Israel) and R. M. Mitterer (Richardson, TX, U.S.A.) 293
Sampling methods for quantitative analysis of coal-ball plants
J. S. Pryor (Athens, OH, U.S.A.) 313
Gradual Late Pliocene onset of glaciation: a deep-sea record from the Northeast Atlantic
P. Loubere (DeKalb, IL, U.S.A.) 327
An isotopic study of a fluvial-lacustrine sequence: the Plio-Pleistocene Koobi Fora sequence, East Africa
T. E. Cerling (Salt Lake City, Utah, U.S.A.), J. R. Bowman (Salt Lake City, Utah, U.S.A.) and J. R. O'Neil (Menlo Park, CA, U.S.A.)..... 335
Biogenic facies in the Antarctic glacimarine environment: basis for a polar glacimarine summary
E. W. Domack (Clinton, NY, U.S.A.) 357